

DATA SET DESCRIPTION

Multi-annual grids of precipitation height over Germany 1971-2000

Version v1.0

Cite data set as: DWD Climate Data Center (CDC): Multi-annual grids of precipitation height over Germany 1971-2000, version v1.0.

INTENT OF THE DATASET

The grids are derived from DWD stations and legally and qualitatively equivalent partner stations in Germany run for climatological and climate related applications, considering the climatology.

POINT OF CONTACT

Deutscher Wetterdienst
CDC - Vertrieb Klima und Umwelt
Frankfurter Straße 135
63067 Offenbach
Tel.: + 49 (0) 69 8062-4400
Fax.: + 49 (0) 69 8062-4499
Mail: klima.vertrieb@dwd.de

DATA DESCRIPTION

Spatial coverage	Germany
Temporal coverage	01.01.1971 - 31.12.2000
Spatial resolution	1 km x 1 km
Temporal resolution	30 years, for each calendar month and season, and for the whole year
Projection	3-degree Gauss-Kruger zone 3, Ellipsoid Bessel, Datum Potsdam (central point Rauenberg), EPSG:31467, see http://spatialreference.org/ref/epsg/31467/ . To define the spatial projection in GIS, the file https://opendata.dwd.de/climate_environment/CDC/help/gk3.prj can be used. Help is given on importing into ESRI ArcGIS in https://opendata.dwd.de/climate_environment/CDC/help/Hilfe_Gauss-Krueger-Raster2GIS.pdf .
Format(s)	There are files for each calendar month (*01.asc.gz bis *12.asc.gz), for each season, i.e., spring (March, April, May): *13.asc.gz, summer (June, July, August): *14.asc.gz, autumn (September, October, November): *15.asc.gz, winter (December, January, February): *16.asc.gz, and for the whole year (*17.asc.gz). The winter value contains the December of the previous year. The file in ESRI-ascii-grid-format has in the header the coordinates for the lower left grid cell, including the definition of its center [XLLCENTER],[YLLCENTER] or its corner [XLLCORNER],[YLLCORNER]. It contains a table of 654 x 866 numbers. Each row goes from West to East. The first row is the northernmost one (654 values with 4 digits). Missing values are marked with -999.
Parameters	Multi-annual mean of precipitation in mm
Uncertainties	Uncertainties are caused by the interpolation method, and erroneous or missing observations. When comparing grid fields for different periods, it should be considered that the measurement network has changed over time.

DATA ORIGIN

The multi-annual means for the calendar months are averages of the respective 30 monthly grids. For the seasonal values (spring-13, summer-14, autumn-15, winter-16) and for the whole year (-17) the respective monthly grids were summed up.

VALIDATION AND UNCERTAINTY ESTIMATE

The given resolution of 1 km x 1 km is the resolution of the employed digital height model. The actual information density depends on the station network.

REFERENCES

Kaspar et al.: Monitoring of climate change in Germany – data, products and services of Germany's National Climate Data Centre. Adv. Sci. Res., 10, 99–106, 2013.

Maier, U. und Müller-Westermeier, G.: Verifikation klimatologischer Rasterfelder, Berichte des Deutschen Wetterdienstes 235, Selbstverlag des Deutschen Wetterdienstes, Offenbach am Main, 2010.

Müller-Westermeier, G., Walter, A., Dittmann, E.: Klimaatlas Bundesrepublik Deutschland, Teil 1-4, Selbstverlag des Deutschen Wetterdienstes, Offenbach am Main, 2005.

Müller-Westermeier, G.: Numerische Verfahren zur Erstellung klimatologischer Karten, Berichte des Deutschen Wetterdienstes 193, Selbstverlag des Deutschen Wetterdienstes, Offenbach am Main, 1995.

COPYRIGHT

The instructions in ftp://ftp-cdc.dwd.de/pub/CDC/Terms_of_use.pdf should be followed. The DWD website provides comprehensive copyright information.

REVISION HISTORY

This document is maintained by DWD division National Climate Monitoring, last edited 18.12.2018.